The maximum number of P_ℓ copies in P_k-free graphs

Ervin Győri — **Alfréd Rényi Institute of Mathematics, Hungarian Academy of Sciences**
Nika Salia — **Central European University**
Casey Tompkins — **Alfréd Rényi Institute of Mathematics, Hungarian Academy of Sciences**
Oscar Zamora — **Central European University**

Abstract

Generalizing Turán's classical extremal problem, Alon and Shikhelman investigated the problem of maximizing the number of T copies in an H-free graph, for a pair of graphs T and H. Whereas Alon and Shikhelman were primarily interested in determining the order of magnitude for large classes of graphs H, we focus on the case when T and H are paths, where we find asymptotic and in some cases exact results. We also consider other structures like stars and the set of cycles of length at least k, where we derive asymptotically sharp estimates. Our results generalize well-known extremal theorems of Erdős and Gallai.

For a graph G, we let $e(G)$ denote the number of edges in G, and for a given graph H, we let $\mathcal{N}(H, G)$ denote the number of (not necessarily induced) copies of H in G. If there is no copy of H in G, we say that G is H-free. We denote the path with k edges by P_k and the cycle with k edges by C_k. By $C_{\geq k}$ we mean the set of all cycles of length at least k. Given a graph G containing a vertex v, we denote the neighborhood of v by $N(v)$. The vertex and edge sets of G are denoted by $V(G)$ and $E(G)$, respectively. Finally, given a set $S \subseteq V(G)$, we denote by $G[S]$ the induced subgraph of G with vertex set S.

Following the notation of Alon and Shikhelman [2], we let $\mathrm{ex}(n, T, H)$ be the maximum number of (noninduced) copies of T in an H-free graph on n vertices. Observe that we have $\mathrm{ex}(n, P_1, H) = \mathrm{ex}(n, H)$, the classical extremal number. If instead of a single graph a set of graphs $\mathcal{H}$ is forbidden, then we define $\mathrm{ex}(n, \mathcal{H})$ (and similarly $\mathrm{ex}(n, T, \mathcal{H})$), in the obvious way.

We begin by recalling the famous theorem of Erdős and Gallai on P_k-free graphs as well as some recent generalizations due to Luo, where the number of cliques is considered.

Theorem 1 (Erdős–Gallai [7])**.** *For all $n \geq k$,*

$$\mathrm{ex}(n, P_k) \leq \frac{(k-1)n}{2},$$

and equality holds if and only if k divides n and G is the disjoint union of cliques of size k.

In their paper, Erdős and Gallai deduced Theorem 1 as a corollary of the following result about graphs with no long cycles.

Theorem 2 (Erdős–Gallai [7])**.** *For all $n \geq k$,*

$$\mathrm{ex}(n, C_{\geq k}) \leq \frac{(k-1)(n-1)}{2},$$

and equality holds if and only if $k-2$ divides $n-1$ and G is a connected graph such that every block is a clique of size $k-1$.




Figure 1: The graph $G_{n,k,a}$ is pictured on the left, and the special case of $G_{n,k,t}$ is pictured on the right. The dashed edge appears only when k is even.

As the extremal examples for Theorem 1 are disconnected, it is natural to consider a version of the problem where the base graph is assumed to be connected. Kopylov [10] settled this problem, and later Ballister, Győri, Lehel and Schelp [3] classified the extremal cases.

We denote by $G_{n,k,a}$ the graph whose vertex set is partitioned into 3 classes, A, B and C with $|A| = a$, $|B| = n - k + a$, $|C| = k - 2a$ such that $A \cup C$ induces a clique, B is an independent set and all possible edges are taken between vertices of A and B.

Throughout this paper we let $t = \lfloor \frac{k-1}{2} \rfloor$. In $G_{n,k,t}$, the class C has one vertex when k is odd or two vertices when k is even. By grouping B and C together, we have that $G_{n,k,t}$ is obtained from a complete bipartite graph $K_{t,n-t}$ by adding all edges in the color class of size t, and in the case that k is even, adding one additional edge inside the color class of size $n - t$.

Theorem 3 (Kopylov [10], Ballister–Győri–Lehel–Schelp [3])**.** *Let G be a connected n-vertex P_k-free graph, with $n \geq k$, then*

$$e(G) \leq \max(e(G_{n,k,t}), e(G_{n,k,1})).$$

Moreover, the extremal graph is either $G_{n,k,t}$ or $G_{n,k,1}$.

Note that if $n \geq 5k/4$, this maximum is achieved by $G_{n,k,t}$. Also observe that

$$e(G_{n,k,t}) = t(n-t) + \binom{t}{2} + \eta_k,$$

where η_k is 1, if k is even, and 0 otherwise. Thus, Theorems 1 and 3 yield the same bound asymptotically as n tends to infinity.

The following theorem was deduced by Luo [11] as a corollary of her main result but also follows from Theorem 1 using a simple induction argument.

Theorem 4 (Luo [11])**.**

$$\mathrm{ex}(n, K_r, P_k) \leq \frac{n}{k}\binom{k}{r}.$$

For our results we will need only that $\mathrm{ex}(n, K_r, P_k) \leq c_{k,r}n$ for some constant $c_{k,r}$ depending only on k and r.

If we impose the additional condition that the graph is connected, then the situation is more complicated. Luo proved the following sharp bounds.

Theorem 5 (Luo [11])**.** *Let $n > k \geq 3$ and G be a connected n-vertex graph with no path of length k, then*

$$\mathcal{N}(K_r, G) \leq \max\left(\mathcal{N}(K_r, G_{n,k,t}), \mathcal{N}(K_r, G_{n,k,1})\right).$$

Theorem 6 (Luo [11]). *Let $n \geq k \geq 4$ and G be a connected n-vertex graph with no cycle of length k or greater, then*

$$\mathcal{N}(K_r, G) \leq \frac{n-1}{k-2}\binom{k-1}{r}.$$

Some recent generalizations of the Erdős–Gallai theorem and Luo's results can be found in [12]. In the present paper we focus on results where paths or all sufficiently long cycles are forbidden. The general problem of enumerating cycles of a fixed length when a fixed cycle is forbidden has also been considered recently (see [8] and [9] which generalize earlier results for special cases, e.g., [4], [5], [2]).

We write $f(n,k,\ell) \sim g(n,k,\ell)$ for $\lim_{k\to\infty}\lim_{n\to\infty} f(n,k,\ell)/g(n,k,\ell) = 1$. (All results for which we use the $\sim$ notation involve parameters n, k and ℓ.) The notation O is always defined in terms of n and may hide constants dependent on k or ℓ.

Alon and Shikhelman [2] considered the problem of maximizing the number of copies of a tree T in a graph which is H-free, for another tree H. Given two trees T and H, they introduced an integer parameter $m(T,H)$ and proved that $\mathrm{ex}(n,T,H) = \Theta(n^{m(T,H)})$, thereby determining the correct order of magnitude for all pairs of trees. In the present paper, we are interested in the case where the forbidden tree is a path, and we find find correct asymptotics and sometimes the exact bound.

We have the following estimates on the number of path and cycle copies in a P_k-free graph.

Theorem 7.

$$\mathrm{ex}(n, P_{2\ell}, P_k) \sim \frac{k^\ell n^{\ell+1}}{2^{\ell+1}}.$$

Theorem 8.

$$\mathrm{ex}(n, P_{2\ell+1}, P_k) \sim \frac{(\ell+2)k^{\ell+1}n^{\ell+1}}{2^{\ell+2}}.$$

Theorem 9.

$$\mathrm{ex}(n, C_{2\ell}, P_k) \sim \frac{k^\ell n^\ell}{\ell 2^{\ell+1}}.$$

Theorem 10.

$$\mathrm{ex}(n, C_{2\ell+1}, P_k) \sim \frac{k^{\ell+1}n^\ell}{2^{\ell+2}}.$$

The construction showing the lower bounds for Theorems 7 through 10 is the same as the extremal construction for the connected version of the Erdős–Gallai theorem, Theorem 3. Because we are interested in asymptotics, we will omit one edge from this construction which only occurs when k is even. Our n-vertex graph G is defined by taking a clique on a set S of $\lfloor \frac{k-1}{2} \rfloor$ vertices and connecting every vertex in S to every vertex of an independent set U, defined on $n - \lfloor \frac{k-1}{2} \rfloor$ vertices. It is easy to see that this graph is P_k-free. In enumerating the $P_{2\ell}$ copies, the only paths which contribute asymptotically alternate between S and U, starting and ending with U (the factor of 2 comes from counting the path in both directions).

When enumerating the $P_{2\ell+1}$ copies, we have two kinds of paths which contribute asymptotically: those that start and end in U, using an edge in S at some step, and those that start in U and end in S, never using an edge contained in S. For the first type, we condition on which step in the path we use the edge in S (ℓ possibilities). Each such path gets counted twice, hence we divide by two. For the second type, each path is counted once and so we do not have to divide by 2.

We present several exact results such as the following.

Theorem 11. *For every positive integer $k \geq 3$, there exists $n_1 \in \mathbb{N}$ such that if $n \geq n_1$*

$$\mathrm{ex}(n, P_2, P_k) = \mathcal{N}(P_2, G_{n,k,t}) = t\binom{n-1}{2} + (n-t)\binom{t}{2} + 2t\eta_k,$$

where $\eta_k = 1$, if k is even, and 0 otherwise. Moreover the only extremal graph is $G_{n,k,t}$.

Theorem 12. *For every positive integer $k \geq 5$, there exists $n_1 \in \mathbb{N}$ such that if $n \geq n_1$*

$$\mathrm{ex}(n, P_3, P_k) = \mathcal{N}(P_3, G_{n,k,t}) = \frac{3t(t-1)}{2}n^2 + O(n).$$

Moreover the only extremal graph is $G_{n,k,t}$.

Theorem 13. *For every positive integer $k \geq 5$, there exists $n_1 \in \mathbb{N}$ such that if $n \geq n_1$*

$$\mathrm{ex}(n, P_4, P_k) = \mathcal{N}(P_4, G_{n,k,t}) = \frac{t(t-1)}{2}n^3 + O(n^2).$$

Moreover the only extremal graph is $G_{n,k,t}$.

References

[1] N. Alon. On the number of subgraphs of prescribed type of graphs with a given number of edges. *Israel Journal of Mathematics*, 38: 116–130, 1981.

[2] N. Alon and C. Shikhelman. Many T copies in H-free graphs. *Journal of Combinatorial Theory, Series B*, 121: 146–172, 2016.

[3] P. N. Balister, E. Győri, J. Lehel, and R. H. Schelp. Connected graphs without long paths. *Discrete Mathematics*, 308(19): 4487–4494, 2008.

[4] B. Bollobás and E. Győri. Pentagons vs. triangles. *Discrete Mathematics,* 308(19): 4332–4336, 2008.

[5] E. Győri and H. Li. The maximum number of triangles in C_{2k+1}-free graphs. *Combinatorics, Probability & Computing,* 21(1–2): 187, 2012.

[6] G. A. Dirac. Some theorems on abstract graphs. *Proceedings of the London Mathematical Society*, 3(1): 69–81, 1952.

[7] P. Erdős and T. Gallai. On maximal paths and circuits of graphs. *Acta Mathematica Hungarica*, 10(3–4): 337–356, 1959.

[8] L. Gishboliner and A. Shapira. A generalized Turán problem and its applications. arXiv preprint arXiv:1712.00831, 2017.

[9] D. Gerbner, E. Győri, A. Methuku, M. Vizer. Generalized Turán problems for even cycles. arXiv preprint arXiv:1712.07079, 2017.

[10] G. N. Kopylov. Maximal paths and cycles in a graph. *DOKLADY AKADEMII NAUK SSSR*, 234(1): 19–21, 1977.

[11] R. Luo. The maximum number of cliques in graphs without long cycles. *Journal of Combinatorial Theory, Series B*, 128: 219–226, 2018.

[12] B. Ning and X. Peng. Extensions of Erdős–Gallai theorem and Luo's theorem with applications. arXiv preprint arXiv:1801.09981, 2018.

[13] V. Nikiforov. The spectral radius of graphs without paths and cycles of specified length. *Linear Algebra and its Applications*, 432(9): 2243–2256, 2010.